ANESTHESIA FOR PATIENTS WITH CARDIOVASCULAR, RESPIRATORY & GASTROINTESTINAL DISEASES

Lyon Lee DVM PhD DACVA DACVECC

Ways to support the cardiovascularly challenged patients

- Stabilize heart rate & rhythm prior to anesthesia if possible
- Optimize cardiac function prior to anesthesia if possible
- Physical examination: observe jugular distension, pulsation, palpate peripheral arterial pulse quality, auscultate heart for assessing characteristics of murmurs
- Thorough cardiac evaluation prior to anesthesia – ECG, Doppler echocardiograph, thoracic radiographs, blood pressure measurement, ultrasonography, cardiac catheterization
- Laboratory evaluation – PCV, TP, hemoglobin content, arterial blood gases, electrolytes
- Choose anesthetic agents that produce minimal cardiovascular changes and preferably have drugs of short duration of action or that are reversible
- Preanesthetics: rely mostly on opioids +/- benzodiazepines: neuroleptanalgesic combination
- Anticholinergics are used judiciously
- Employ local anesthetic technique under sedation or even general anesthesia
- Induction: propofol, etomidate, ketamine, mask with inhalant
- Maintenance: usually isoflurane or sevoflurane (rapid recovery and less cardiovascular depression than halothane)
- Monitor cardiovascular performance
  - ECG: rate and rhythm
  - arterial blood pressure (BP = CO x PVR)
  -
    - x

General considerations

- Many anesthetics produce some degree of respiratory depression
- The respiratory depression in combination with cardiovascular depression induced by

- Propofol
- Etomidate
- Ketamine

- minimize oxygen deficit period by allowing rapid intubation and ventilation
- Control airway as quickly as possible, begin positive pressure ventilation (esp. with lower airway disease)
- Nitrous oxide may be better avoided.
  - It diffuses into gaseous pocket and worsens symptoms such as pneumothorax
  - It reduces the inspiratory fraction of oxygen
- Monitoring:
  - ECG
  - Pulse oximetry
  - BP
  - Capnography
  - Serial blood gas analysis
  - Tidal volume and peak airway pressure (thoracic compliance)
  - Temperature
- Recovery
  - Maintain ET tube in situ as long as possible
  - Post-operative pulse oximetry
  - Support ventilation as long as possible
  - Consider post anesthetic oxygen supplementation
    - mask
    - nasal catheter
    - oxygen cage
  -

Case example 2

- "Miss Genuines"
- Signalment: 1 week old Quarter Horse filly
- History: presented for choanal atresia
- Significant physical exam findings: normal neonatal foal except for nasal obstruction
- Laboratory finding: normal
- Referring DVM had performed a tracheostomy shortly after birth
- Presented for anesthesia 4/6 for laser surgical correction of choanal atresia

Preanesthetic management?

Anesthetic induction?

Maintenance of anesthesia?

Monitoring?

Postoperative care?

General considerations

- Variety of disease processes...
- Malabsoption
- Derangement of electrolytes, acid-base status
- hypovolemia
-

Equine Colic

- One of our most common emergency surgical procedures
- Patients present in a variety of conditions, from minimally to severely compromised
  - Respiratory compromise
  - Cardiovascular compromise
  - Dehydration
  - Hypotension
  - Hypoxemia
  - Electrolyte imbalances
  - Acid/Base disturbances
- Again, stabilize if possible
  - Large volumes of fluids IV rapidly (multiple large bore catheters)
  - Bicarbonate if acidotic
  - Pain management (usually w/ alpha-2, NSAID such as Flunixin meglumine)
- Our current anesthetic protocol