

PREANESTHETIC EVALUATIONS & ECG

Lyon Lee DVM PhD DACVA

"If you don't look, you don't see"

PREANESTHETIC EVALUATIONS & PREPARATIONS

Introduction

- The preanesthetic examination or evaluation will influence greatly on the dose and choice of the premedicants, induction agents and maintenance agents as well as selection of anesthetic techniques.
- Thorough patient evaluation and preparation will improve patient safety and ensure successful anesthetic outcome.

Preparation of the patient

Food withholding time

- Varies between species
- Dogs and cats : usually 12 hours and minimum of 6 hours
-

Venous access

- Placement of intravenous catheter facilitate IV administration of induction agents, and also minimize the extravascular injection of irritant agents such as thiopental sodium.
- For dogs and cats 20 - 22 G, 1 - 1.5 inch long are most commonly used, and for large animals 12 – 14 G of 3 - 4 inch long are most commonly used.

Items you need to prepare on a tray for a cat presented for general anesthesia at the BVMTH


- Three ET tubes (one that would fit best, each of smaller and larger)
- Eye lube (ophthalmic ointment for eye lubrication)
- Gauze roll bandage
- One inch and half inch tapes
- Stylet
- Two to three heparinized saline flush in 3 ml syringe
- Laryngoscope and blades (size 1 and 2)
- Cuff syringe
- 4 x 4 gauze pad
- K-Y jelly
- Lidocaine 2 % 0.25 ml in TB syringe
- Two catheters 20 - 22 G, 1 – 1.5 inch long
- Injection cap
- Needles of varying size (20 G x 1”, 22 G x 1 “)
-

- Remember there is no safe anesthetic agent but only safe anesthetist, so ensure every effort to minimize overall risk based on your evaluation and plan.

ELECTROCARDIOGRAPHY

Electrocardiogram

- ECG is recommended for dogs and cats over 7 years old prior to general anesthesia as part of preanesthetic work-up to screen underlying systemic diseases and abnormalities.
- Graphic recording of electrical potentials produced by cardiac muscle during different phases of the cardiac cycle



ECG paper chart

- @ 25 mm/sec
 - one small square; 0.04 sec
 - five small squares; 0.2 sec
- @ 50 mm/sec
 - one small square; 0.04 sec
 - five small squares; 0.2 sec

Five physiologic properties of the heart

- Automaticity
- Excitability

- Record a brief tracing at 50 mm/sec for ease of assessment of P-QRS-T waveforms
- Observe the following during the ECG recording
 - Whether the top and bottom of the waveform are all seen
 - Adjust the alignment as appropriate
- Decrease the sensitivity to $\frac{1}{2}$ cm = 1 mV if QRS complexes go off the paper
- Increase the length of the trace if arrhythmia is present
- R waves should be positive in lead I if negative, check the lead wires to determine whether they are attached to the correct limbs. If correct, then a true abnormality exists.

Canine ECG normal values

Rate

- 70 to 160 beats/ min for adult dogs

•

T wave

- Can be positive, negative, or diphasic
- Not greater than $\frac{1}{4}$ amplitude of R wave

Q-T interval

- Width: 0.15 to 0.25 sec ($7\frac{1}{4}$ - $12\frac{1}{4}$ boxes) at normal heart rate

Feline ECG Normal Values

Rate

- 120 to 240 beats/mm

Rhythm

- Normal sinus rhythm
- Sinus tachycardia (physiologic reaction to excitement)

P wave

- Width: maximum, 0.04 sec (2 boxes wide)
- Height: maximum, 0.2 mV (2 boxes tall)

P-R interval

- Width: 0.05 to 0.09 sec ($2\frac{1}{4}$ to $4\frac{1}{4}$ boxes)

QRS complex

- Width: maximum, 0.04 sec (2 boxes)
- Height of R wave: maximum, 0.9 mV (9 boxes)

S-T segment

- No depression or elevation

T wave

- Can be positive, negative, or diphasic; most often positive
- Maximum amplitude: 0.3 mV (3 boxes)

- Third-degree atrioventricular block
- Left bundle branch block
- Right bundle branch block

Interpreting Arrhythmias —the Simple Steps

- Arrhythmias can be intimidating. Therefore, it is important that we find a simple approach for analyzing rhythm strips. Systematically following the five-step method outlined below has proven to be both simple and effective.

Step 1. Calculate the heart rate

- Decide whether the heart rate is rapid, slow, or normal.

Step 2. Assess the rhythm

- Scan the strip from left to right, noting if the R-R intervals are regular or irregular.
- A caliper is a handy tool for plotting P-P and R-R intervals.

R intervals —

- P waves may precede normal QRS complexes by different time spans.
- **Long P-R intervals**—indicate an AV conduction delay
- (1° AV block) Td ()Tj EMC /LBody <</MCID 10 4>BDC /C2_0 1 Tf -237.481